

		References
1.	SCHEDULING STATUS	
2.	S3	
3.	1. NAME OF THE MEDICINE	
4.	DIACOMIT 250 Capsules	
5.	DIACOMIT 500 Capsules	
6.	DIACOMIT 250 Powder for oral suspension	
7.	DIACOMIT 500 Powder for oral suspension	
8.	2. QUALITATIVE AND QUANTITATIVE COMPOSITION	
9.	DIACOMIT 250 Capsules: each hard capsule contains 250 mg stiripentol.	
10.	DIACOMIT 500 Capsules: each hard capsule contains 500 mg stiripentol.	
11.	DIACOMIT 250 Powder for oral suspension: each sachet contains 250 mg stiripentol as the	
12.	active ingredient.	
13.	DIACOMIT 500 Powder for oral suspension: each sachet contains 500 mg stiripentol as the	
14.	active ingredient.	
15.	DIACOMIT is sugar free.	
16.	For the full list of excipients, see section 6.1.	
17.	3. PHARMACEUTICAL FORM	
18.	Hard capsules.	
19.	Powder for oral suspension.	

20. DIACOMIT 250 Capsules: opaque pink capsules of size 2 with “Diacomit” printed in
21. black on cap and “250 mg” printed in black on body with self-locking closure containing a
22. white to pale yellow powder.
23. DIACOMIT 500 Capsules: opaque white capsules of size 0 elongated (0+) with “Diacomit”
24. printed in black on cap and “500 mg” printed in black on body with self-locking closure
25. containing a white to pale yellow powder.
26. DIACOMIT 250 and 500 Powder for oral suspension is a pale pink powder filled in a
27. single dose sachet.
28. **4. CLINICAL PARTICULARS**
29. **4.1 Therapeutic indications**
30. DIACOMIT is indicated for adjunctive treatment of generalised tonic-clonic and clonic
31. seizures associated with severe myoclonic epilepsy in infancy (SMEI, also known as Dravet
32. syndrome) in patients whose seizures are not adequately controlled with a benzodiazepine
33. (usually clobazam) and valproate.
34. **4.2 Posology and method of administration**
35. **Posology**
36. Prescriptions and treatment should be initiated by neurologists experienced in the diagnosis
37. and management of epilepsy, with continuation of patient management by general
38. paediatricians and practitioners when the initiating neurologist is unavailable.
39. **General**
40. The dose of DIACOMIT is calculated on a mg/kg body weight basis.

41. It is recommended to split the daily dose in two or three daily intakes (totalling the daily
 42. recommended dose per kg and per day). The initiation of adjunctive therapy with
 43. DIACOMIT should be undertaken gradually using upwards dose escalation to reach the
 44. recommended dose of 50 mg/kg/day.

45. DIACOMIT dosage escalation should be gradual, starting with 20 mg/kg/day for 1 week,
 46. then 30 mg/kg/day for 1 week. Further dosage escalation is age dependent:
 47. - children less than 6 years should receive an additional 20 mg/kg/day in the third week,
 48. thus achieving the recommended dose of 50 mg/kg/day in three weeks;
 49. - children from 6 to less than 12 years should receive an additional 10 mg/kg/day each
 50. week, thus achieving the recommended dose of 50 mg/kg/day in four weeks;
 51. - children and adolescents 12 years and older should receive an additional 5 mg/kg/day
 52. each week until the optimum dose is reached based on clinical judgement.

53. The recommended dose of 50 mg/kg/day is based on the available clinical study findings
 54. and was the only dose of DIACOMIT evaluated in the pivotal studies (see Clinical Trials
 55. subheading under section 5.1).

56. There are no clinical study data to support the clinical safety of DIACOMIT administered
 57. at daily doses greater than 50 mg/kg/day.

58. There are no clinical study data to support the use of DIACOMIT as monotherapy in Dravet
 59. syndrome.

60. **Dose adjustments of other antiepileptics used in combination with DIACOMIT**

61. Despite the absence of comprehensive pharmacology data on potential drug interactions, the
 62. following advice regarding modification of the dose and dosage schedules of other
 63. antiepileptic medicines administered in conjunction with DIACOMIT is provided based on
 64. clinical experience.

65. *Clobazam*

66. In the pivotal studies, when the use of stiripentol was initiated, the daily dose of clobazam
 67. was 0,5 mg/kg/day usually administered in divided doses, twice daily. In the event of clinical
 68. signs of adverse reactions or overdose of clobazam (i.e., drowsiness, hypotonia, and
 69. irritability in young children), this daily dose was reduced by 25 % every week.
 70. Approximately two to three-fold increases in clobazam and fivefold increases in
 71. norclobazam plasma levels respectively have been reported with co-administration of
 72. DIACOMIT in children with Dravet syndrome.

73. *Valproate*

74. The potential for metabolic interaction between DIACOMIT and valproate is considered
 75. modest and thus, no modification of valproate dosage should be needed when DIACOMIT
 76. is added, except for clinical safety reasons. In the pivotal studies in the event of
 77. gastrointestinal adverse reactions such as loss of appetite, loss of weight, the daily dose of
 78. valproate was reduced by around 30 % every week.

79. **Effect of formulation**

80. The sachet formulation ($C_{\max} = 7,32 \mu\text{g/ml}$) has a slightly higher $C_{\max}$ than the capsules ($C_{\max}$
 81. $= 5,99 \mu\text{g/ml}$) and thus the formulations are not bioequivalent. It is recommended that if a
 82. switch of formulations is required this is done under clinical supervision, in case of problems
 83. with tolerability (see section 5).

84.	Renal and hepatic impairment
85.	DIACOMIT is not recommended for use in patients with impaired hepatic and/or renal
86.	function (see section 4.4).
87.	Method of administration
88.	The capsule should be swallowed whole with a glass of water during a meal.
89.	The powder should be mixed in a glass of water and should be taken immediately after
90.	mixing during a meal.
91.	DIACOMIT must always be taken with food as it degrades rapidly in an acidic environment
92.	(e.g. exposure to gastric acid in an empty stomach).
93.	DIACOMIT should not be taken with milk or dairy products (yoghurt, soft cream cheese,
94.	etc.), carbonated drinks, fruit juice or food and drinks that contain caffeine or theophylline.
95.	Discontinuation of DIACOMIT
96.	Antiepileptic medicines, including DIACOMIT, should be withdrawn gradually to minimise
97.	the potential for seizures or increased seizure frequency. In clinical trials in children with
98.	Dravet syndrome, dosages were gradually reduced over a period lasting at least one month.
99.	In a situation where rapid withdrawal of DIACOMIT is medically required, appropriate
100.	monitoring is recommended.
101.	4.3 Contraindications

102.	Hypersensitivity to stiripentol or to any of the excipients of DIACOMIT as listed in section
103.	6.1
104.	A past history of psychoses in the form of episodes of delirium.
105.	4.4 Special warnings and precautions for use
106.	<i>Identified precautions</i>
107.	The pivotal clinical evaluation of DIACOMIT was in children of 3 years of age and over
108.	with SMEI. Children under 3 years of age and adults were not studied in the pivotal trials
109.	therefore efficacy and safety for those populations were not established in the pivotal trials.
110.	Medicine interactions
111.	<i>Carbamazepine, phenytoin and phenobarbital</i>
112.	Carbamazepine, phenytoin, and phenobarbital should generally not be used in conjunction
113.	with DIACOMIT in the management of Dravet syndrome.
114.	Phenytoin and phenobarbital can significantly decrease the serum concentration of
115.	stiripentol. DIACOMIT can significantly increase the serum concentration of
116.	carbamazepine, phenytoin, and phenobarbital. Furthermore, there is evidence that these
117.	medicines have the potential to worsen seizure activity in patients with Dravet syndrome,
118.	even though these medicines may have a role in a subset of patients with Dravet syndrome.
119.	It is recommended that carbamazepine, phenytoin, and phenobarbital be used in conjunction
120.	with DIACOMIT only under expert supervision with appropriate monitoring.
121.	<i>Clobazam and valproate</i>

122. The daily dosage of clobazam and/or valproate should be reduced according to the onset of
 123. side effects whilst on DIACOMIT therapy (see section 4.2).

124. *Substances interfering with CYP enzymes*

125. Stiripentol as contained in DIACOMIT is an inhibitor of the enzymes CYP1A2, CYP2C8,
 126. CYP2C19, CYP3A4, CYP2C9 and CYP2D6 and may markedly increase the plasma
 127. concentrations of substances metabolised by these enzymes and increase the risk of adverse
 128. reactions (see section 4.5). *In vitro* data also suggest that STP induces the activity of
 129. CYP1A2 and CYP2B/3A, but the clinical relevance of this is uncertain. *In vitro* studies
 130. suggested that stiripentol phase 1 metabolism is catalysed by CYP1A2, CYP2C9, CYP2C19
 131. and CYP3A4/5 and possibly other enzymes. Caution is advised when combining stiripentol
 132. with other substances that inhibit or induce one or more of these enzymes.

133. **Blood count**

134. Neutropenia may be associated with the administration of DIACOMIT, clobazam and
 135. valproate. Blood counts should be assessed prior to starting treatment with DIACOMIT.
 136. Unless otherwise clinically indicated, blood counts should be checked every 6 months.

137. **Neurologic/psychiatric**

138. Patients should be monitored for somnolence and drowsiness, particularly when
 139. DIACOMIT is used concomitantly with other central nervous depressant. An adjustment of
 140. the dosage of or other anti-epileptic medicines could be considered. Patients (or their parents
 141. or carers) should be advised to not operate machinery or drive.

142. Movement disorders including ataxia, hypotonia, tremor hyperkinesia, dysarthria and
 143. equilibrium disorders have been reported in patients treated with DIACOMIT for Dravet
 144. syndrome.

145. ***Suicidal ideation and behaviour***

146. Suicidal ideation and behaviour have been reported in patients treated with antiepileptic
 147. medicines in several indications.

148. All patients treated with antiepileptic medicines, irrespective of indication, should be
 149. monitored for signs of suicidal ideation and behaviour and appropriate treatment should be
 150. considered. Patients (or their parents or carers) should be advised to seek medical advice
 151. should signs of suicidal ideation or behaviour emerge.

152. ***Use in hepatic impairment***

153. **Liver function**

154. In patients newly prescribed DIACOMIT, raised γ GT (notably when combined with
 155. valproate) are observed with a common frequency, whilst abnormal liver function tests
 156. (increases in AST and/or ALT) are also reported with a rare frequency. Liver function (γ GT,
 157. AST and ALT levels) should therefore be assessed prior to starting treatment with
 158. DIACOMIT and, unless otherwise clinically indicated, should be checked every 6 months.

159. Stiripentol as contained in DIACOMIT is primarily metabolised by the liver and primarily
 160. excreted by the kidney.

161.	In the absence of specific clinical data in patients with impaired hepatic or renal function,
162.	DIACOMIT is not recommended for use in patients with impaired hepatic and/or renal
163.	function.
164.	<i>Use in renal impairment</i>
165.	Stiripentol as contained in DIACOMIT is primarily metabolised by the liver and primarily
166.	excreted by the kidney.
167.	In the absence of specific clinical data in patients with impaired hepatic or renal function,
168.	DIACOMIT is not recommended for use in patients with impaired hepatic and/or renal
169.	function.
170.	<i>Use in the elderly</i>
171.	No data are available.
172.	<i>Paediatric use</i>
173.	Growth rate of children
174.	Due to the frequency of gastrointestinal adverse reactions to treatment with DIACOMIT and
175.	valproate (anorexia, loss of appetite, nausea, vomiting), the growth rate of children under
176.	this combination of treatment should be carefully monitored.
177.	Effects on laboratory tests
178.	In the event of an abnormal blood count or liver function test finding, the clinical decision
179.	for continuing use or adjusting the dose of DIACOMIT in conjunction with adjusting the
180.	doses of clobazam and valproate needs to be made on an individual patient basis taking into
181.	consideration the potential clinical benefits and risks.

182.	4.5 Interaction with other medicines and other forms of interaction	
183.	Stiripentol as contained in DIACOMIT is subject to non-linear pharmacokinetics (see	
184.	section).	
185.	Potential medicine interactions affecting DIACOMIT	
186.	The influence of other antiepileptic medicines on stiripentol pharmacokinetics is not well	
187.	established.	
188.	The impact of macrolides and azole antifungal medicines, that are known to be inhibitors of	
189.	CYP3A4 and substrates of the same enzyme, on stiripentol metabolism is not known.	
190.	Likewise, the effect of DIACOMIT on their metabolism is not known.	
191.	<i>In vitro</i> studies suggested that stiripentol phase 1 metabolism is catalysed by CYP1A2,	
192.	CYP2C19 and CYP3A4 and possibly other enzymes. Caution is advised when combining	
193.	DIACOMIT with other substances that inhibit or induce one or more of these enzymes.	
194.	Effect of DIACOMIT on cytochrome P450 enzymes	
195.	Many of these interactions have been partially confirmed by <i>in vitro</i> studies and in clinical	
196.	trials. The increase in steady state levels with the combined use of stiripentol, valproate, and	
197.	clobazam is similar in adults and children, though inter-individual variability is marked.	
198.	At therapeutic concentrations, stiripentol significantly inhibits several CYP450 isoenzymes:	
199.	for example, CYP2C19, CYP1A2, CYP2C8, CYP2D6, CYP2C9 and CYP3A4. As a result,	
200.	pharmacokinetic interactions of metabolic origin with other medicines may be expected.	
201.	These interactions may result in increased systemic levels of these active substances that	

202. may lead to enhanced pharmacological effects and to an increase in adverse reactions. *In*
 203. *vitro* data also suggest that stiripentol induces the activity of CYP1A2 and CYP2B/3A, but
 204. the clinical relevance of this is uncertain.

205. Caution must be exercised if clinical circumstances require combining stiripentol with
 206. CYP2C19 (e.g. citalopram, omeprazole) or CYP3A4 (e.g. HIV protease inhibitors,
 207. antihistamines such as astemizole and chlorpheniramine, calcium channel blockers, statins,
 208. oral contraceptives, codeine) due to the increased risk of adverse reactions (see further in
 209. this section for antiepileptic medicines). Monitoring of plasma concentrations or adverse
 210. reactions is recommended. A dose adjustment may be necessary.

211. Co-administration with CYP3A4 substrates with a narrow therapeutic index should be
 212. avoided due to the markedly increased risk of severe adverse reactions.

213. Data on the potential for inhibition of CYP1A2 are limited, and therefore, interactions with
 214. theophylline and caffeine cannot be excluded because of the increased plasma levels of
 215. theophylline and caffeine which may occur via inhibition of their hepatic metabolism,
 216. potentially leading to toxicity. Use in combination with DIACOMIT is not recommended.
 217. This warning is not only restricted to medicines but also to a considerable number of foods
 218. and nutritional products aimed at children, such as cola drinks, which contain significant
 219. quantities of caffeine or chocolate, which contains trace amounts of theophylline.

220. As stiripentol inhibited CYP2D6 *in vitro* at concentrations that are achieved clinically in
 221. plasma, substances that are metabolised by this isoenzyme like: beta-blockers (propranolol,
 222. carvedilol, timolol), antidepressants (fluoxetine, paroxetine, sertraline, imipramine,
 223. clomipramine), antipsychotics (haloperidol), analgesics (codeine, dextromethorphan,

tramadol) may be subject to metabolic interactions with DIACOMIT. A dose-adjustment may be necessary for substances metabolised by CYP2D6 and that are individually dose titrated.

Effect of DIACOMIT on GABA receptor site

Stiripentol is a positive allosteric modulator of the GABA_A receptor. There is a potential for it to act synergistically with other allosteric modulators at the same site (e.g. benzodiazepines, non-benzodiazepines, barbiturates, bromides or neuroactive steroids) to enhance GABAergic neurotransmission.

Potential for DIACOMIT to interact with other medicines

In the absence of available clinical data, caution should be taken with the following clinically relevant interactions with DIACOMIT:

Drug or drug class	Clinical comment
<i>Undesirable combinations (to be avoided unless strictly necessary)</i>	
Rye ergot alkaloids (ergotamine, dihydroergotamine)	Ergotism with possibility of necrosis of the extremities (inhibition of hepatic elimination of rye ergot).
Cisapride, halofantrine, pimozone, quinidine, bepridil	Increased risk of cardiac arrhythmias and torsades de pointes/wave burst dysrhythmia in particular.
Immunosuppressants (tacrolimus, ciclosporin, sirolimus)	Raised blood levels of immunosuppressants (decreased hepatic metabolism).
Statins (atorvastatin, simvastatin, etc.)	Increased risk of dose-dependent adverse reactions such as rhabdomyolysis (decreased hepatic metabolism of cholesterol-lowering

248.		medicine).
249.	<i>Combinations requiring precautions</i>	
250.	Midazolam, triazolam, alprazolam	Increased plasma benzodiazepine levels may
251.		occur via decreased hepatic metabolism
252.		leading to excessive sedation.
253.	Chlorpromazine	Enhanced central depressant effect of
254.		chlorpromazine.
255.	Effects on other Anti-epileptic medicines (AEDs)	
256.	Concomitant antiepileptic	Clinical comment
257.	medicine	
258.	Phenobarbitone,	Inhibition of CYP450 isoenzyme CYP2C19 and CYP3A4
259.	carbamazepine, phenytoin,	may provoke pharmacokinetic interactions (inhibition of
260.	primidone, clobazam,	their hepatic metabolism) with phenobarbitone, primidone,
261.	valproate, diazepam,	phenytoin, carbamazepine, clobazam, valproate (refer to
262.	ethosuximide, and tiagabine.	section 4.4), diazepam (enhanced myorelaxation),
263.		ethosuximide, and tiagabine. The consequences are
264.		increased plasma levels of these anticonvulsants with
265.		potential risk of overdose.
266.		
267.		Phenytoin, phenobarbitone, carbamazepine should not be
268.		used in conjunction with DIACOMIT in Dravet patients.
269.		
270.		Clinical monitoring of plasma levels of other
271.		anticonvulsants when combined with DIACOMIT with
272.		possible dose adjustments is recommended.

273.	Topiramate	In a compassionate use program for stiripentol, topiramate
274.		was added to stiripentol, clobazam and valproate in 41 %
275.		of 230 cases. Based on the clinical observations in this
276.		group of patients, there is no evidence to suggest that a
277.		change in topiramate dose and dosage schedules is needed
278.		if co-administered with stiripentol.
279.		
280.		With regard to topiramate, it is considered that potential
281.		competition of inhibition on CYP2C19 should not occur
282.		because it probably requires plasma concentrations 5-15
283.	times higher than plasma concentrations obtained with the	
284.	standard recommended topiramate dose and dosage	
285.	schedules.	
286.	Levetiracetam	Levetiracetam does not undergo hepatic metabolism to a
287.		major extent. As a result, no pharmacokinetic metabolic
288.		drug interaction between stiripentol and levetiracetam is
289.		anticipated.
290.	CYP3A4 inhibitors, such a DIACOMIT, may increase the exposure of oestrogen or	
291.	progestin containing contraceptives when used concomitantly. Oestrogen or progestin	
292.	containing contraceptives may increase the exposure of CYP1A2 or CYP3A substrates, such	
293.	a stiripentol. The clinical significance of these potential interactions is not known.	
294.	4.6 Fertility, pregnancy and lactation	
295.	Women of childbearing potential / Contraception in males and females	

296. The use of efficient methods of contraception is advisable.

297. **Pregnancy**

298. *Risk related to epilepsy and antiepileptic medicines in general.*

299. It has been shown that in the offspring of women with epilepsy, the prevalence of
 300. malformations is two to three times greater than reported in the general population. Although
 301. other factors, e.g. the epilepsy, can contribute, available evidence suggests that this increase,
 302. to a large extent, is caused by the treatment. In the treated population, an increase in
 303. malformations has been noted with polytherapy.

304. However, effective anti-epileptic therapy should not be interrupted during pregnancy, since
 305. the aggravation of the illness may be detrimental to both the mother and the foetus.

306. *Risk related to stiripentol*

307. No data on exposed pregnancies are available. DIACOMIT and/or its metabolites cross the
 308. placenta in rats. There was no evidence of teratogenicity in mice or rabbits treated with STP
 309. during the period of organogenesis at oral doses up to 800 mg/kg/day, which is about 2-fold
 310. (mice) and 6-fold (rabbits) the MRHD, based on body surface area. Increased numbers of
 311. resorptions were reported in mouse embryofoetal development studies at doses of ≥ 200
 312. mg/kg/day (0,4 times the MRHD based on BSA), and also in the rabbit embryofoetal
 313. development study at a dose of 800 mg/kg/day (6 times the MRHD based on BSA). Rats
 314. treated with oral STP from early or late gestation to weaning showed maternotoxicity and
 315. reduced pup survival and reflex development at 800 mg/kg/day (equivalent to about 3-fold
 316. the mg/m² MRHD); the no-effect dose was 200 mg/kg/day.

317. In view of the indication, administration of DIACOMIT during pregnancy and in women of
 318. childbearing potential would not be expected. The clinical decision for use of DIACOMIT
 319. in pregnancy needs to be made on an individual patient basis.

320. **Breastfeeding**

321. There are no human studies on the excretion of DIACOMIT in breast milk. DIACOMIT
 322. passes freely from plasma into milk in the goat. Women taking DIACOMIT should not
 323. breastfeed their infants as adverse events are expected.

324. **Fertility**

325. There were no effects on fertility in male and female rats following oral administration of
 326. STP at doses up to 800 mg/kg/day (equivalent to about 3-fold the MRHD, based on body
 327. surface area). As no human clinical data are available, the potential risk for humans is
 328. unknown.

329. **4.7 Effects on ability to drive and use machines**

330. Patients with Dravet syndrome would not be expected to drive or operate machinery due to
 331. the nature of the underlying disease and the effects of long-term administration of
 332. anticonvulsant medicines.

333. DIACOMIT may cause dizziness and ataxia that may affect ability to drive and use
 334. machines and patients should not drive or use machinery whilst on DIACOMIT therapy.

335. **4.8 Undesirable effects**

336. *a. Summary of the safety profile*

337. Two prospective, randomised, 8-week treatment, double-blind, placebo-controlled trials
 338. were conducted in Dravet syndrome patients (STICLO France and STICLO Italy). In both
 339. trials, stiripentol was added on to treatment with valproate and clobazam. The dose of
 340. stiripentol was 50 mg/kg/day.

341. The following table lists the adverse events that were reported during the studies. No patient
 342. died during these double-blind, placebo-controlled studies.

343. **Table 1 Number and percentage of patients with Dravet syndrome who experienced**
 344. **adverse events in double-blind placebo-controlled clinical studies in which stiripentol**
 345. **(STP) was used as an adjunct treatment with valproate and clobazam**

Body System / Adverse event	STICLO Pooled Total	
	STP N = 33	Placebo N = 31
Number of Patients with at Least One Adverse event	31 (94 %)	12 (39 %)
Body as a Whole – General Disorders		
Asthenia/fatigue	2 (6 %)	-
Central and Peripheral Nervous System Disorders		
Drowsiness/sleepiness	22 (67 %)	3 (10 %)
Ataxia	4 (12 %)	3 (10 %)
Hypotonia	5 (15 %)	1 (3 %)
Tremor	3 (9 %)	1 (3 %)
Hyperkinesia	(3 %)	2 (6 %)
Dysarthria	2 (6 %)	-
Equilibrium disorders	1 (3 %)	-
<i>Status epilepticus</i>	1 (3 %)	1 (3 %)
Motor deficiency	-	1 (3 %)
Gastrointestinal System Disorders		
Loss of appetite	13 (39 %)	2 (6 %)
Weight loss	8 (24 %)	-
Nausea/vomiting	5 (15 %)	1 (3 %)
Sialorrhea	2 (6 %)	-
Body System / Adverse event	STICLO Pooled Total	
	STP N = 33	Placebo N = 31
Weight gain	5 (15 %)	4 (13 %)
Abdominal pain	3 (9 %)	1 (3 %)
Diarrhoea	-	1 (3 %)

Laboratory Parameters		
Neutropenia	3 (9 %)	-
Thrombocytopenia	2 (6 %)	-
Increase in aspartate aminotransferase	1 (3 %)	-
Eosinophilia	1 (3%)	-
Psychiatric Disorders		
Hyperexcitability/agitation	7 (21 %)	1 (3 %)
Aggressiveness/irritability	5 (15 %)	1 (3 %)
Insomnia/nightmares	2 (6 %)	-
Intellectual slowing	1 (3 %)	-
Respiratory System Disorders		
Bronchitis	1 (3 %)	1 (3 %)
Rhinitis	1 (3 %)	1 (3 %)
Skin and Appendages Disorders		
Face erythema	1 (3 %)	-
Dry skin	1 (3 %)	-
Urticaria	1 (3 %)	-
Urinary System Disorders		
Dysuria	1 (3 %)	-

346. **Gastrointestinal adverse events**
347. *Weight loss and anorexia*
348. Anorexia and weight loss are very common adverse effects.
349. In two double-blind, placebo-controlled trials in Dravet syndrome patients, gastrointestinal
350. adverse events most often reported in patients receiving stiripentol were loss of appetite in
351. 39 % of the treated patients and loss of weight in 24 % of the treated patients.
352. *Nausea and vomiting*
353. Nausea and vomiting are common adverse effects.
354. In 2 double-blind, placebo-controlled trials in Dravet syndrome patients, nausea and
355. vomiting were observed in 15 % of the treated patients.

356. Due to the frequency of gastrointestinal adverse reactions to treatment with stiripentol and
 357. valproate (anorexia, loss of appetite, nausea, vomiting), the growth rate of children under
 358. this combination of treatment should be carefully monitored.

359. *Abnormal liver function tests*

360. Raised γ GT (notably when combined with carbamazepine and valproate) is a common
 361. adverse effect.

362. Liver function should be assessed prior to starting treatment with DIACOMIT. Unless
 363. otherwise clinically indicated, liver function should be checked every 6 months. As the
 364. medicine is metabolised mainly by the liver, DIACOMIT is not recommended for use in
 365. patients with impaired hepatic function.

366. **Central nervous system adverse events**

367. Drowsiness, ataxia, hypotonia, dystonia and insomnia are very common adverse effects.

368. In 2 double-blind, placebo-controlled trials in Dravet syndrome patients,
 369. drowsiness/sleepiness were observed in 67 % of the treated patients, hyperexcitability/
 370. agitation in 21 % of treated patients, aggressiveness/irritability in 15 % of treated patients,
 371. hypotonia in 15% of the treated patients, and ataxia in 12 % of the treated patients.

372. Patients should be monitored for somnolence and drowsiness, particularly when
 373. DIACOMIT is used concomitantly with other central nervous depressant. An adjustment of
 374. the dosage of concomitant clobazam or other anti-epileptic medicines could be considered.
 375. Patients should be advised to not operate machinery or drive.

Thrombocytopenia and neutropenia

Neutropenia is a common adverse effect. Thrombocytopenia is a rare adverse effect.

In 2 double-blind, placebo-controlled trials in Dravet syndrome patients, neutropenia was observed in 9 % of the treated patients and thrombocytopenia in 6 % of the treated patients.

Blood counts should be assessed prior to starting treatment with DIACOMIT. Unless otherwise clinically indicated, blood counts should be checked every 6 months.

Post marketing experience

The most frequent [side effects with DIACOMIT are anorexia, weight loss, insomnia, drowsiness, ataxia, hypotonia and dystonia.

Table 2 Tabulated list of DIACOMIT Adverse Effects

System Class (MedDRA)	Organ	Frequent	Frequency unknown
Blood and lymphatic system disorder		Neutropenia, persistent severe neutropenia usually resolves spontaneously when DIACOMIT is stopped*.	Thrombocytopenia*
Metabolism and nutrition disorders		Anorexia*, loss of appetite*, weight loss (especially when combined with sodium valproate)*	
Psychiatric disorder		Insomnia*, aggressiveness*, irritability*, behaviour	

399.		disorders*, opposing	
400.		behaviour*,	
401.		hyperexcitability*, sleep	
402.		disorders*	
403.	Nervous system	Drowsiness*, ataxia*,	
404.	disorders	hypotonia*, dystonia*,	
405.		hyperkinesias	
406.			
407.	Eye disorders		Diplopia (when used in
408.			combination with
409.			carbamazepine)
410.	Gastrointestinal	Nausea*, vomiting*	
411.	disorders		
412.	Skin and		Photosensitivity, rash,
413.	subcutaneous tissue		cutaneous allergy, urticaria
414.	disorders		
415.	General disorders		Fatigue
416.	and admini-		
417.	stration site		
418.	conditions		
419.	Investigations	Raised γ GT (notably when	Liver function test abnormal
420.		combined with carbamaze-	
421.		pine and valproate)	
422.	<u>Infections and</u>		<u>Pneumonia and aspiration</u>
423.	<u>infestations</u>		<u>pneumonia</u>

Ref. 1, pg.
11, 4.8.9

424.	Many of the above adverse reactions are often due to an increase in plasma levels of other	
425.	anticonvulsant medicines (see section 4.4 and section 4.5) and may regress when the dose	
426.	of these medicines is reduced.	
427.	* Data are derived from both clinical trials and post-marketing experience.	
428.	Reporting of suspected adverse reactions	
429.	Reporting suspected adverse reactions after authorisation of the medicine is important. It	Type IA_{IN} Code C.I.0.2a Editorial change to update the requirements for reporting of adverse events. As per SAHPRA PI guideline, SAHPGL-CEM-02_v6
430.	allows continued monitoring of the benefit/risk balance of the medicine. Healthcare	
431.	providers are asked to report any suspected adverse reactions to SAHPRA via the [“6.04	
432.	Adverse Drug Reaction Reporting Form” , found online under SAHPRA’s publications:	
433.	https://www.sahpra.org.za/Publications/Index/8. Med Safety APP (Medsafety X	
434.	SAHPRA) and eReporting platform (who-umc.org) found on SAHPRA website.	
435.	<u>For reporting of adverse reactions directly to the Holder of Certificate of Registration, email:</u>	
436.	<u>pharmacovigilance@equitypharma.co.za.</u>	
437.	4.9 Overdose	
438.	Data on clinical overdose are not available. Treatment is supportive (symptomatic measures	
439.	in intensive care units).	
440.	5. PHARMACOLOGICAL PROPERTIES	
441.	5.1 Pharmacodynamic properties	
442.	Pharmacotherapeutic group: Antiepileptics, other antiepileptics, ATC code: N03AX17	
443.	Mechanism of action	

444. The mechanism of antiepileptic activity of stiripentol is based on the potentiation of the
445. GABAergic transmissions in the CNS. *In vitro*, stiripentol has been shown to directly
446. enhance GABAA receptor-mediated transmission by acting both post-synaptically at a
447. neuronal site coupled to the GABAA receptor and pre-synaptically to increase GABA
448. release from nerve terminals. In rodent models, stiripentol appears to increase brain levels
449. of GABA. This could occur by inhibition of synaptosomal uptake of GABA and/or
450. inhibition of GABA transaminase. Stiripentol has been shown to enhance GABAA receptor-
451. mediated transmission in the immature rat hippocampus and increase the mean open-
452. duration (but not the open-frequency) of GABAA receptor chloride channels.

453. Stiripentol also potentiates the efficacy of clobazam and other anticonvulsants, as a result of
454. the pharmacokinetic interactions (see section 4.5). This effect of stiripentol is mainly based
455. on metabolic inhibition of several isoenzymes, in particular CYP450 3A4 and 2C19,
456. involved in the hepatic metabolism of other anti-epileptic medicines.

457. **Clinical Trials**

458. The efficacy of stiripentol, as an add-on therapy to optimised treatment with valproate and
459. clobazam in patients presenting with Dravet syndrome has been demonstrated in two
460. randomised, double-blind, placebo-controlled clinical trials. These trials, designated
461. STICLO France and STICLO Italy, involved 41 (age range: 3,0-20,7) and 23 patients (age
462. range: 3,5- 18,9) respectively. Both studies were conducted according to similar design to
463. evaluate the responder rate to treatment, where the responder was defined as a patient who
464. achieved ≥ 50 % decrease in the frequency of generalised clonic or tonic-clonic seizures
465. during the double-blind treatment period compared to baseline (i.e., placebo run-in).

466.	Eligible patients were randomly allocated to receive either stiripentol (50 mg/kg/day) or	
467.	placebo added on to their antiepileptic treatment.	
468.	In study STICLO France, total of 41 patients were enrolled; 21 were randomised to receive	
469.	stiripentol and 20 were randomised to receive placebo. One patient in the stiripentol group	
470.	was excluded. For study STICLO Italy, 23 patients were enrolled; 12 were randomised to	
471.	STP, and 11 to placebo. All were judged to be evaluable. A summary of the baseline	
472.	demographic data is provided in Table 3.	
473.		Ref 1, p. 14 (5.1.7)
		Type IA_{IN} Code C.I.0.2a Editorial update - “The reference citation ‘Ref 1, p.14 (5.1.7) was unintentionally included in the PI during document and has been deleted.

Table 3 Baseline Demographic and Illness Characteristics across the Pivotal Trials STICLO France and STICLO Italy						
Baseline characteristics	STICLO France N=41		STICLO Italy N=23		STICLO Pooled Total N=64	
	STP N=21	Placebo N=20	STP N=12	Placebo N=11	STP N=33	Placebo N=31
Gender						
Male	6 (29 %)	11 (55 %)	8 (67 %)	5 (46 %)	14 (42 %)	16 (52 %)
Female	15 (71 %)	9 (45 %)	4 (33 %)	6 (54 %)	19 (58 %)	15 (48 %)
Age (years)						
Mean ± SD	9,4 ± 4,0	9,3 ± 4,9	9,2 ± 3,6	8,7 ± 4,4	9,3 ± 3,8	9,1 ± 4,6
Median	9,8	9,2	8,6	8,2	8,6	8,8
Min-Max	3,0 – 16,7	3,2 – 20,7	3,7 – 15,5	3,5 – 18,9	3,0 – 16,7	3,2 – 20,7
Age Group (n, %)						
< 1	0 (0 %)	0 (0 %)	0 (0 %)	0 (0 %)	0 (0 %)	0 (0 %)
1 to < 6	4 (19 %)	6 (30 %)	3 (25 %)	3 (27 %)	7 (21 %)	9 (29 %)
6 to < 12	10 (48 %)	9 (45 %)	6 (50 %)	6 (54 %)	16 (48 %)	15 (48 %)
12 to < 17	7 (33 %)	2 (10 %)	3 (25 %)	1 (9 %)	10 (30 %)	3 (10 %)
≥ 17	0 (0 %)	3 (15 %)	0 (0 %)	1 (9 %)	0 (0 %)	4 (13 %)
Weight (kg)						
n	20	20	12	11	32	31
Mean ± SD	31,8 ± 12,7	30,5 ± 14,4	31,9 ± 11,7	29,2 ± 9,0	31,8 ± 12,1	30,0 ± 12,6
Median	30,0	25,5	29,2	27,0	30,0	27,0
Min - Max	14 - 60	15 – 70	16 -55	18 - 49	14 - 60	15 - 70
SCN1A mutation (n, %)*						

n	20	Not tested	9	Not tested	29	Not tested
Mutation	14 (70 %)		9 (100 %)		23 (79,3 %)	
No mutation	6 (30 %)		0 (0 %)		6 (20,7 %)	
Disease Duration (years)						
Mean ± SD	9,1 ± 3,8	9,0 ± 5,0	9,1 ± 3,6	8,7 ± 4,5	9,1 ± 3,7	8,9 ± 4,7
Min - Max	3,0 - 16,7	3,2 - 20,7	3,7 - 15,2	3,2 - 18,9	3,0 - 16,7	3,2 - 20,7
Seizure type (n, %)						
Generalised Seizures						
Tonic-clonic	21 (100 %)	20 (100 %)	12 (100 %)	11 (100 %)	33 (100 %)	31 (100 %)
Myoclonic	10 (47,6 %)	14 (70,0 %)	10 (83,3 %)	8 (72,7 %)	20 (60,6 %)	22 (71,0 %)
Unspecified generalised seizure	12 (57,1 %)	12 (60,0 %)	5 (41,7 %)	5 (45,5 %)	17 (51,5 %)	17 (54,8 %)
Number of tonic-clonic or clonic seizures during baseline						
Mean ± SD	17,9 ± 17,3	18,5 ± 17,0	33,6 ± 28,2	27,5 ± 28,7	23,6 ± 22,8	27,1 ± 21,8
Median	11,4	14,4	30,2	20,7	14,5	15,5
Min - Max	4 - 73	4 - 76	2 - 86	4 - 101	2 - 86	4 - 101
Daily dose Clobazam (mg/kg/day)						
n	21	20	12	11	33	31
Mean ± SD	0,5 ± 0,2	0,5 ± 0,2	0,7 ± 0,2	0,5 ± 0,1	0,6 ± 0,2	0,5 ± 0,2
Median	0,5	0,5	0,6	0,5	0,5	0,5
Min - Max	0 - 1	0 - 1	0 - 1	0 - 1	0 - 1	0 - 1
Daily dose Valproate (mg/kg/day)						
n	21	20	12	11	33	31
Mean ± SD	21,7 ± 9,6	20,7 ± 8,3	29,8 ± 7,8	25,6 ± 7,3	24,7 ± 9,7	22,4 ± 8,2
Median	17,0	16,7	29,1	26,9	21,9	22,5
Min - Max	11 - 43	14 - 44	21 - 44	11 - 38	11 - 44	11 - 44

474. STP = Stiripentol SD = Standard deviation
475. *: the presence or absence of the SCN1A mutation was measured after the blind was
476. broken and the presence or absence of the mutation was only studied in the STP groups.
477. **Table 4 summarises the antiepileptic efficacy of stiripentol in each STICLO trial, as**
478. **well as the results of the pooled analysis (“STICLO Total”).**

Table 4 STICLO Efficacy Results Obtained at the End of Treatment in the Intent-to-Treat Population across the Pivotal Trials STICLO France and STICLO Italy						
	STICLO France N=41		STICLO Italy N=23		STICLO Pooled Total N=64	
	STP	Placebo	STP	Placebo	STP	Placebo
Percentage change from baseline in seizure frequency*						
n	21	20	11	9	32	29
Mean ± SD	-62,0 ±	12,1 ± 44,4	-74 ± 26,7	-13 ± 62,0	-66 ± 44,2	4,3 ± 50,7
Median	-87,5	12,1	-81,2	-27,4	-84,4	-5,8
Min-Max	-100 – 72,6	-75 – 119	-100 – -33	-87 – 140	-100 – 72,6	-87 – 140
p-value ¹	0,0003		0,0056		<0,0001	
Responder analysis†						
No of	15/21	1/20	8/12	1/11	23/33	2/31

responders/total						
(Responder rate)	(71,4 %)	(5,0 %)	(66,7 %)	(9,1 %)	(69,7 %)	(6,5 %)
[95 %CI]	[52,1 –	[0,0 – 14,6]	[40,0 –	[0,0 – 26,1]	[54,0 –	[0,0 – 15,1]
p-value ²	< 0,0001		0,0098		<0,0001	

479. *: Frequency of generalised tonic-clonic or clonic seizures.

480. †: Responder is defined as a patient with a $\geq 50\%$ decrease in frequency of generalised
481. tonic-clonic or clonic seizures.

482. [1] Wilcoxon Test; [2] Fisher's Exact Test.

483. CI=confidence interval; SD=standard deviation.

484. When efficacy results of both STICLO trials were pooled, 23 of 33 (69,7 %) patients on
485. stiripentol versus 2 of 31 (6,5 %) patients on placebo met the criterion for response ($p <$
486. 0,0001). Mean percent reduction in seizure frequency was 66 % in the pooled STP group
487. versus an increase of 4 % in the pooled placebo groups ($p < 0,0001$).

488. 5.2 Pharmacokinetic properties

489. Absorption

490. The T_{max} times were 2,42 hours for single doses of stiripentol 500 mg and 1 000 mg and
491. 2,96 hours for 2 000 mg. According to Michaelis-Menten kinetics there is some evidence
492. for non-linearity regarding stiripentol absorption. Following the oral administration of 500
493. mg, 1 000 mg and 2 000 mg of stiripentol in healthy volunteers the following data were
494. determined.

495.

	500 mg	1 000 mg	2 000 mg
C_{max} (mg/L)			
Mean $\pm$ SD	2,63 $\pm$ 1,18	6,63 $\pm$ 1,83	13,8 $\pm$ 4,83
Median	2,14	6,50	14,1
Min-max	1,21 – 4,1	3,90 – 10,4	8,31 – 24,0
AUC _{0-30 h} (mg/L.h)			
Mean $\pm$ SD	8,85 $\pm$ 3,77	32,1 $\pm$ 10,7	79,0 $\pm$ 24,2

Median	8,13	30,2	82,0
Min-max	3,74 – 15,7	18,3 – 51,1	48,2 - 128

The absolute bioavailability of stiripentol is not known since an intravenous formulation is not available for testing. It is well absorbed by the oral route since the majority of an oral dose is excreted in urine. Relative bioavailability between the capsules and powder for oral suspension in sachet formulations has been studied in healthy male volunteers after a 1 000 mg single oral administration. $C_{\max}$ of the sachet was slightly higher (23 %) compared with the capsule and did not meet the criteria for bioequivalence. $T_{\max}$ was similar with both formulations. Clinical supervision is recommended if switching between the stiripentol capsule and powder for oral suspension in sachet formulations.

Distribution

Stiripentol binds extensively to circulating plasma proteins (about 99 % at clinical plasma concentrations).

In a population pharmacokinetic study conducted in Dravet syndrome patients at steady state receiving the combination valproate + clobazam + stiripentol, the apparent volume of stiripentol distribution according to weight was as follows.

Body weight (kg)	V/F (L)
10	32,0 ± 3,8
30	95,9 ± 11,5
60	191,8 ± 23,0

Biotransformation

Stiripentol is primarily metabolised by the liver.

Stiripentol undergoes extensive first-pass metabolism in the liver with CYP1A2,

514. CYP2C19 and CYP3A4 being the main isozymes involved in stiripentol metabolism.
 515. Stiripentol is mainly metabolised by glucuronidation and oxidative cleavage.
 516. Stiripentol is supplied as a racemic mixture, and after administration R-enantiomer is the
 517. predominant enantiomer, but glucuronidation appears to favour the S-enantiomer.
 518. Elimination of a single dose was mainly (73 %) via the kidney after extensive
 519. metabolization (13 different metabolites) by the liver (Cytochrome P 450).

520. **Elimination**

521. Stiripentol is primarily excreted by the kidney.
 522. Systemic exposure to stiripentol increases markedly compared to dose proportionality. In
 523. healthy adults, plasma clearance decreases markedly at high doses; it falls from
 524. approximately 40 l/kg/day at the dose of 600 mg/day to about 8 l/kg/day at the dose of
 525. 2,40 mg. Clearance is decreased after repeated administration of stiripentol, probably due
 526. to inhibition of the cytochrome P450 isoenzymes responsible for its metabolism. The half-
 527. life of elimination was in the range of 4,5 hours to 13 hours, increasing with dose.
 528. Similarly, in a population pharmacokinetic study conducted in Dravet syndrome patients
 529. at steady state receiving the combination valproate + clobazam + stiripentol, the apparent
 530. volume of stiripentol distribution according to weight was as follows.

531.

Body weight (kg)	CL/F (L/h)	T _½ (h)
10	2,60 ± 0,18	8,5 ± 1,3
30	4,19 ± 0,29	15,9 ± 2,4
60	5,65 ± 0,40	23,5 ± 3,5

532. **5.3 Preclinical safety data**

533. **Genotoxicity**

534. There was no evidence for genotoxic potential in *in vitro* assays for gene mutation or
 535. unscheduled DNA synthesis. A positive *in vitro* chromosomal aberration assay in Chinese
 536. Hamster Ovary cells (observed at the highest concentration) was not confirmed in a similar
 537. assay with human lymphocytes or in a mouse micronucleus test *in vivo*. The genotoxic
 538. potential of STP is considered to be low.

539. **Carcinogenicity**

540. There was no evidence of carcinogenicity in rats following oral administration of STP for
 541. 2 years at doses up to 800 mg/kg/day. This dose is approximately 3-4 times the MRHD
 542. based on body surface area, and systemic exposure (plasma AUC) was about 4-fold the
 543. estimated exposure in adults at the MRHD. In mice treated orally for 78 weeks, there was
 544. an increase in the incidence of hepatic adenomas and carcinomas at doses of 200 or 600
 545. mg/kg/day. These doses are about 0,5 to 2-fold the MRHD based on body surface area,
 546. and plasma exposure was about 2-fold the paediatric C_{max}; the no-effect dose was 60
 547. mg/kg/day. In view of the lack of genotoxicity of STP and the known susceptibility of the
 548. mouse liver to tumour formation in response to hepatic enzyme induction, this finding is
 549. not considered to indicate a risk of tumorigenicity in patients.

550. **6 PHARMACEUTICAL PARTICULARS**

551. **6.1 List of excipients**

552. *Capsule:*

553. Magnesium stearate

554. Povidone

555. Sodium starch glycolate, type A

556. *Capsule shell (Body & cap):*

557.	Erythrosine (E127)	
558.	Gelatin	
559.	Indigotin carmine (E132)	
560.	Titanium dioxide (E171)	
561.	Powder for oral suspension:	
562.	Aspartame	
563.	Spray-dried liquid glucose	
564.	Povidone	
565.	Sodium starch glycolate type A	
566.	Erythrosine	
567.	Titanium dioxide	
568.	Carmellose	
569.	Hyetellose	
570.	Arome Polv Tutti Frutti 25 H 245	
571.	6.2 Incompatibilities	
572.	Not applicable.	
573.	6.3 Shelf life	
574.	3 years.	
575.	6.4 Special precautions for storage	
576.	Store at or below 25 °C.	
577.	Store in the original container to protect from light.	
578.	6.5 Nature and contents of container	

579.	Diacomit Capsules	
580.	DIACOMIT 250 and 500 mg capsules are packaged in opaque polyethylene bottles closed	
581.	with a child-resistant tamper-evident polypropylene screw cap.	
582.	The DIACOMIT 250 mg capsules are packaged in 60 mL bottles and the DIACOMIT 500	
583.	mg capsules in 100 mL bottles in cardboard cartons.	
584.	Each bottle of 250 or 500 mg capsules contains 60 capsules.	
585.	Diacomit Powder for oral suspension	
586.	Diacomit 250 & 500 Powder for oral suspensions are packaged in sachets consisting of a	
587.	paper, aluminium foil and polyethylene film composite. The sachets supplied in cardboard	
588.	boxes with 30, 60 or 90 sachets.	
589.	6.6 Special precautions for disposal and other handling	
590.	Any unused medicine or waste material should be disposed of in accordance with local	
591.	requirements.	
592.	7. HOLDER OF CERTIFICATE OF REGISTRATION	
593.	Equity Pharmaceuticals (Pty) Ltd	
594.	100 Sovereign Drive	
595.	Route 21 Corporate Park	
596.	Nellmapius Drive	
597.	Irene, 0157	
598.	Pretoria	
599.	<u>Tel: +27 (0) 12 345 1747</u>	Type IA_{IN} Code C.I.0.2a Editorial update to include the telephone number
600.	8. REGISTRATION NUMBER(S)	

601.	Diacomit 250 Capsule: 550281	
602.	Diacomit 500 Capsule: 550282	
603.	Diacomit 250 Powder for oral suspension: 550283	
604.	Diacomit 500 Powder for oral suspension: 550284	
605.	9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION	
606.	07 June 2022	
607.	10. DATE OF REVISION OF THE TEXT	
608.	To be allocated by SAHPRA.	

References:

Reference 1	Approved TGA Innovator PI, DIACOMIT, Chiesi Australia Pty Ltd, date of revision of text: 25 August 2025, pages 1-20	Module 1.3.1.2.2